INDEX, VOLUME 49, 1956

A-42, for control of cucumber beetles, 132
Abundance of beet leafhopper, factors affecting, 95
Acalymma vittata, 131, 384
Acanthocinus obsoletus, 664
Acarina, 219
Acarus siro, 813
Aceratogallia sanguinolenta, 371
Aceria tulipae, 1, 405
Acheta domestica, 262
Acontia dacia, 764
Acricidae, 371
Acrobasis caryae, 404
Adalia bipunctata, 384
Adistemia watsoni, 269
Adjuvants, for use with polyhedrosis viruses, 703
Aedes
  aegypti, 156, 313, 564, 610
  albopictus, 261
  dorsalis, 87
  pseudoscutellaris, 786
  sollicitans, 416
  stimulans, 416
  vexans, 416
Agallia constricta, 371
Agria affinis, 413
Agrotis
  gladiaria, 403
  orthogonia, 145
  ypsilon, 403
Agyrotaenia velutinana, 196
Ahasverus advena, 119
Aircraft, control of forest insects by, 92
Air drift of mites, 624
Alabama argillacea, 239, 706, 757, 783
Aldrin, effect of on
  corn root growth, 230
  germination and emergence of sugar beets, 710
Aldrin, for control of
  Acontia dacia, 765
  billbugs on corn, 234
  cabbage maggot, 355
  clover leaf weevils, 543
  clover root borer, 162, 641
  cotton insects, 72
  corn earworm, 540
  corn rootworm, 709
  cucumber beetles, 131
  Drosophila, 709
  dusky sap beetle, 540
  European corn borer, 835
  fowl tick, 449
  green June beetle, 829, 869
  house fly larvae, 75
  meadow spittlebug, 162
  onion thrips, 334
  peanut insects, 69
  potato flea beetle, 531
  rice weevil, 64
  southern potato wireworm, 879
  sweet potato insects in soil 767
  Tabanus larvae, 402
  thief ant, 329
  thrips on peanuts, 632
  tobacco wireworm, 345
  wireworms, 66, 112, 480
Aldrin, toxicity of, to
  boll weevil, 758
  Typhlodromus fallacis, 513
Allethrin
  crystalline isomer of, as a standard of comparison for fly sprays, 450
  effect on acetone on toxicity of, 849
  effectiveness with synergists against the body louse, 644
  for control of *Drosophila*, 609
  formulations in a thermal fog machine, 610
  radioactive, contaminants in, 135
Alphitobius diaperinus, 155
Amblyomerus verditer, 413
Amblyomma americanum, 88, 393
Am. Cyanamid 3911
  effect of on germination and emergence of sugar beets, 710
Am. Cyanamid 3911, for control of apple leaf miner, 218
  clover leaf weevils, 543
  green peach aphid, 26
  European corn borer, 835
Am. Cyanamid 4124, for control of house fly, 27, 78, 622, 705, 752
  northern fowl mite, 348
Am. Cyanamid 12008, for control of cotton insects, 73
  green peach aphid, 26
  southern garden leafhopper, 358
Am. Cyanamid 12009, for control of cotton insects, 73
  southern garden leafhopper, 358
Amphimallon majalis, 179
Anagrus
  armatus nigriceps, 438
  epos, 438
Anastrepha
  ludens, 178, 570
  mombinpraeoptans, 177, 715
Anaiis quindecimpunctata, 384
Anopheles
  crucians, 273
  hyrcanus sinensis, 261
  punctipennis, 273
  quadrimaculatus, 156, 433, 523, 567
Anrhomyza sp., 419
Anthomyiidae, 248
Anthonomus grandis, 72, 239, 696, 706, 712, 757, 783, 874, 877
Anthrenus
  flavipes, 277
  scrophulariae, 277
  verbasci, 277
Antibiosis properties of sorghums against corn leaf aphid, 504
Anuraphis roseus, 522
Aonidiella
  aurantii, 103, 534
  citrina, 534
Apanteles
  anganleti, 268
  diacrisiae, 119
  glomeratus, 321
Apechthis ontario, 413
Aphaereta sp. near *auripes*, 462
Aphrophora parallela, 426
Aphids, low-elevation movement of, 771
Aphis
  abbreviata, 252, 772
  gossypii, 72, 698, 706, 712, 760, 783
  pomi, 522
Apis mellifera, 254, 666, 735, 741, 743, 854
Apis mellifera, queens whose eggs fail to hatch, 11
Aplomya caesar, 413
Araecerus fasciculatus, 119
Aramite, for control of
  cotton insects, 72
  desert spider mite, 860
  European red mite, 350
  mites in earthworm beds, 410
  two-spotted spider mite, 151, 243, 350
  winter grain mite, 519
Aramite, toxicity tests for control of
  desert spider mite, 762
  house cricket, 262
  two-spotted spider mite, 762
  Typhlodromus fallacis, 513
Argas persicus, 447, 713
Arilus cristatus, 384
Aroclor, use of, in emulsifiable concentrates, 719
Aromatic naptha, 132, 415
Arsenical compounds, for control of
  boll weevil, 241
  cotton leafworm, 241
  rice weevil, 241
  southern armyworm, 241
Arsenical compounds, organic, evaluation of, 239
Arsenic tolerance, differences in, among cockroaches, 39
Asperigillus, 67
Asidiotus forbesi, 200
Aspilota, 419
Atrichopogon levis, 419
Attagenus piceus, 154, 277, 724
Aulacizes irrorata, 384
Aulocara elliotti, 618
Autographa californica, 403
Azobenzene, for control of two-spotted spider mite, 151

Bacillus
  larvae, 470
  popilliae, 4
Bacillus thuringiensis, for control of lepidoptera on crucifers, 320
Bacterium, to control lepidopterous pests of crucifers, 320
Baits
  for trapping insects associated with oak wilt, 224, 226
  sprays for *Drosophila* control, 708
Baits, for control of
  flies in Wyoming, 343
  green June beetle larvae, 868
  horn fly, 27
  house fly, 27, 77, 751
  stable fly, 28
Barium fluosilicate, for control of green June beetle larvae, 868
Bathyplectes curculionis, 443
Bayer L 13/59, effect of
  against grain mite, 813
  piperonyl butoxide on anticholinesterase activities on house fly ChE, 591
Bayer L 13/59, for control of
  boll weevil, 713, 784
  bollworm, 713, 784
  cotton aphid, 713
  Drosophila, 709
  flies in Wyoming, 343
  green June beetle, 829
  green peach aphid, 25
  house fly, 27, 78, 705, 752

INDEX, VOLUME 49, 1956

A-42, for control of cucumber beetles, 132
Abundance of beet leafhopper, factors affecting, 95
Acalymma vittata, 131, 384
Acanthocinus obsoletus, 664
Acarina, 219
Acarus siro, 813
Aceratogallia sanguinolenta, 371
Aceria tulipae, 1, 405
Acheta domestica, 262
Acontia dacia, 764
Acricidae, 371
Acrobasis caryae, 404
Adalia bipunctata, 384
Adistemia watsoni, 269
Adjuvants, for use with polyhedrosis viruses, 703
Aedes
  aegypti, 156, 313, 564, 610
  albopictus, 261
  dorsalis, 87
  pseudoscutellaris, 786
  sollicitans, 416
  stimulans, 416
  vexans, 416
Agallia constricta, 371
Agria affinis, 413
Agrotis
  gladiaria, 403
  orthogonia, 145
  ypsilon, 403
Agyrotaenia velutinana, 196
Ahasverus advena, 119
Aircraft, control of forest insects by, 92
Air drift of mites, 624
Alabama argillacea, 239, 706, 757, 783
Aldrin, effect of on
  corn root growth, 230
  germination and emergence of sugar beets, 710
Aldrin, for control of
  Acontia dacia, 765
  billbugs on corn, 234
  cabbage maggot, 355
  clover leaf weevils, 543
  clover root borer, 162, 641
  cotton insects, 72
  corn earworm, 540
  corn rootworm, 709
  cucumber beetles, 131
  Drosophila, 709
  dusky sap beetle, 540
  European corn borer, 835
  fowl tick, 449
  green June beetle, 829, 869
  house fly larvae, 75
  meadow spittlebug, 162
  onion thrips, 334
  peanut insects, 69
  potato flea beetle, 531
  rice weevil, 64
  southern potato wireworm, 879
  sweet potato insects in soil 767
  Tabanus larvae, 402
  thief ant, 329
  thrips on peanuts, 632
  tobacco wireworm, 345
  wireworms, 66, 112, 480
Aldrin, toxicity of, to
  boll weevil, 758
  Typhlodromus fallacis, 513
Allethrin
  crystalline isomer of, as a standard of comparison for fly sprays, 450
  effect on acetone on toxicity of, 849
  effectiveness with synergists against the body louse, 644
  for control of *Drosophila*, 609
  formulations in a thermal fog machine, 610
  radioactive, contaminants in, 135
Alphitobius diaperinus, 155
Amblyomerus verditer, 413
Amblyomma americanum, 88, 393
Am. Cyanamid 3911
  effect of on germination and emergence of sugar beets, 710
Am. Cyanamid 3911, for control of apple leaf miner, 218
  clover leaf weevils, 543
  green peach aphid, 26
  European corn borer, 835
Am. Cyanamid 4124, for control of house fly, 27, 78, 622, 705, 752
  northern fowl mite, 348
Am. Cyanamid 12008, for control of cotton insects, 73
  green peach aphid, 26
  southern garden leafhopper, 358
Am. Cyanamid 12009, for control of cotton insects, 73
  southern garden leafhopper, 358
Amphimallon majalis, 179
Anagrus
  armatus nigriceps, 438
  epos, 438
Anastrepha
  ludens, 178, 570
  mombinpraeoptans, 177, 715
Anaiis quindecimpunctata, 384
Anopheles
  crucians, 273
  hyrcanus sinensis, 261
  punctipennis, 273
  quadrimaculatus, 156, 433, 523, 567
Anrhomyza sp., 419
Anthomyiidae, 248
Anthonomus grandis, 72, 239, 696, 706, 712, 757, 783, 874, 877
Anthrenus
  flavipes, 277
  scrophulariae, 277
  verbasci, 277
Antibiosis properties of sorghums against corn leaf aphid, 504
Anuraphis roseus, 522
Aonidiella
  aurantii, 103, 534
  citrina, 534
Apanteles
  anganleti, 268
  diacrisiae, 119
  glomeratus, 321
Apechthis ontario, 413
Aphaereta sp. near *auripes*, 462
Aphrophora parallela, 426
Aphids, low-elevation movement of, 771
Aphis
  abbreviata, 252, 772
  gossypii, 72, 698, 706, 712, 760, 783
  pomi, 522
Apis mellifera, 254, 666, 735, 741, 743, 854
Apis mellifera, queens whose eggs fail to hatch, 11
Aplomya caesar, 413
Araecerus fasciculatus, 119
Aramite, for control of
  cotton insects, 72
  desert spider mite, 860
  European red mite, 350
  mites in earthworm beds, 410
  two-spotted spider mite, 151, 243, 350
  winter grain mite, 519
Aramite, toxicity tests for control of
  desert spider mite, 762
  house cricket, 262
  two-spotted spider mite, 762
  Typhlodromus fallacis, 513
Argas persicus, 447, 713
Arilus cristatus, 384
Aroclor, use of, in emulsifiable concentrates, 719
Aromatic naptha, 132, 415
Arsenical compounds, for control of
  boll weevil, 241
  cotton leafworm, 241
  rice weevil, 241
  southern armyworm, 241
Arsenical compounds, organic, evaluation of, 239
Arsenic tolerance, differences in, among cockroaches, 39
Asperigillus, 67
Asidiotus forbesi, 200
Aspilota, 419
Atrichopogon levis, 419
Attagenus piceus, 154, 277, 724
Aulacizes irrorata, 384
Aulocara elliotti, 618
Autographa californica, 403
Azobenzene, for control of two-spotted spider mite, 151

Bacillus
  larvae, 470
  popilliae, 4
Bacillus thuringiensis, for control of lepidoptera on crucifers, 320
Bacterium, to control lepidopterous pests of crucifers, 320
Baits
  for trapping insects associated with oak wilt, 224, 226
  sprays for *Drosophila* control, 708
Baits, for control of
  flies in Wyoming, 343
  green June beetle larvae, 868
  horn fly, 27
  house fly, 27, 77, 751
  stable fly, 28
Barium fluosilicate, for control of green June beetle larvae, 868
Bathyplectes curculionis, 443
Bayer L 13/59, effect of
  against grain mite, 813
  piperonyl butoxide on anticholinesterase activities on house fly ChE, 591
Bayer L 13/59, for control of
  boll weevil, 713, 784
  bollworm, 713, 784
  cotton aphid, 713
  Drosophila, 709
  flies in Wyoming, 343
  green June beetle, 829
  green peach aphid, 25
  house fly, 27, 78, 705, 752

northern fowl mite, 348
pink bollworm, 560
Bayer L 13/59, P^{32}-labeled
absorption and translocation of, in a cockroach, 808
metabolism of, in a cow, 801
Bayer L 13/59, resistance of house fly larvae to, 463
Bayer L 13/59, toxicity of to
corn earworm larvae, 409
tobacco budworm larvae, 409
Bayer 16259, for control of
boll weevil, 784
bollworm, 784, 874
pink bollworm, 874
Bayer 17147, for control of
Acontia dacia, 765
boll weevil, 713, 784, 874
bollworm, 713, 784, 874
cabbage looper, 386
cotton aphid, 707, 713
cotton leafworm, 707
green peach aphid, 26
pink bollworm, 874
Bayer 17147, toxicity of to
boll weevil, 759
corn earworm larvae, 409
tobacco budworm larvae, 409
Bayer 21/199, for control of
house fly, 622
northern fowl mite, 348
Bayer 21/199, resistance of house fly larvae to, 463
Bdellonyssus sylviarum, 347
Beauveria globulifera, 441
Bee-collected vs. hand collected apple pollen, 827
Beet yellows virus, transmission of, 789
Bembecia marginata, 287
Bembidion quadramaculatum, 337
Bembix, 203
Benzal azine, toxicity of, to *Typhlodromus fallacis*, 513
Bergoldia virulenta, for control of lepidopterous pests on crucifers, 320
BHC, effect of on corn root growth, 231
BHC, for control of
Acontia dacia, 765
alfalfa weevil, 637
apple leaf miner, 218
boll weevil, 697, 784
bollworm, 784
clover root borer, 162, 641
clover root curculio, 641
corn rootworms, 709
cotton insects, 72
green June beetle larvae, 868
meadow spittlebug, 162, 637, 641
onion thrips, 334
pea aphid, 637
peanut insects, 69
rice weevil, 64
Tabanus larvae, 402
taxus mealybug, 274
tomato russet mite, 769
BHC, toxicity of to
boll weevil, 758
cotton aphid, 762
sheep, 89
BHC, translocation of from soil treatments, 815
Bibio xanthopus, 336
Biology of
a Cimbicid sawfly, *Zaraea americana*, 701
Bdella depressa, 745
black flies, 121
boll weevil, 877
Chrysopid predators of cotton aphid, 698
Drosophila on tomatoes, 607
European corn borer in Alabama, 411
Ferrisiana virgata on azaleas, 276
filth flies in Manitoba, 595
geminate leafhopper, 272
insects associated with oak wilt, 219
Lathridiidae, 269
legume mite, 291
lesser peach tree borer, 397
lupine maggot, 248
Lygus hesperus, 690
pea aphid, 878
peach tree borer, 397
pickleworm, 870
pine sawfly, effect of different hosts on, 400
pink bollworm, 562, 876
Pleocoma oregonensis, 431
Pseudococcus cuspidatae, 273
spiny rat mite, 702
Tachysphex spp., as predators of a grasshopper, 615
winter grain mite, 515
Biological control of
apple pests, 196
fruit flies, 176
hymenopterous predators as agents, 202
Biological races or corn leaf aphid, 499
Biting fly control, effect of, on milk production, 465
Blattella germanica, 130, 411, 662, 714, 820
Blissus insularis, 83
Bordeaux, for control of potato flea beetle, 530
Bourletiella hortensis, 745
Boricola
borisi, 138, 239
caprae, 88
BR-5, 5%, for control of cattle lice, 237
Brachmia rufescens, 205
Brachyacantha ursina, 384
Brachymeria ovata, 413
Bracon
brevicornis, 268
hebetor, 119, 629
Bradysia sp., 621
Brevicoryne brassicae, 328
Brevipalpus inornatus, 392
Bryobia praetiosa, 201, 745
Bucculatrix thurberiella, 174
Bulan, for control of
beet leafhopper, 124
cucumber beetles, 131
Butoxy polypropylene glycol, for control of,
biting flies, 465
Drosophila, 418, 609
horn fly, 396, 407
house fly, 28
stable fly, 396

Calcium arsenate, for control of,
cotton leafworm, 759
Drosophila, 609
potato flea beetle, 530
Calendra
callosa, 233
maidis, 233
Caloglyphus anomalus, 410
Callisto geminatella, 217
Callitroga hominivorax, 86
Caloglyphus spinitarsus, 584
Cantharis rectus, 384
Capitophorus fragaefolii, 243
Captan, for control of apple pests, 523
Captan, harmless to *Typlodromus*, 201
Captan, toxicity of, to *Typhlodromus fallacis*, 512
Carcelia lagoae, 39
Carcinops sp., 155
Carpocapsa pomonella, 196, 467, 522, 866
Carpophilus
dimidiatus, 119
lugubris, 222, 539
Carriers, DDT-impregnated, effect of different types on European corn borer control, 836
Cartodere
constricta, 269
filiformis, 269
Cathartus quadricollis, 119, 753
Cattle Insecticide 5%, for control of cattle lice, 237
Cephalonomia tarsalis, 119
Ceratitis capitata, 134, 135, 176
Ceratomegilla fuscilabris, 384
Cerceris rybyensis, 203
Cercyon sp., 337
Cerotoma trifurcata, 384
Chaetocnema confinis, 766
Chaoborus astictopus, 206
Chelonus
heliopae, 268
n. sp., 268
Chilicola, 203
Chironomidae, 208
Chlordane, effect of on
germination and emergence of sugar beets, 710
yields of red clover, 372
Chlordane, for control of
Adistemia watsoni, 270
banana weevils, 297
billbugs on corn, 234
brown dog tick, 282
cattle lice, 139, 237
chinch bug, 84
clover leaf weevils, 543
clover root borer, 162, 741
cucumber beetles, 132
German cockroach, 130
green June beetle, 829, 869
house cricket, 262
Japanese beetle, 566
house fly larvae, 75
meadow spittlebug, 162
onion thrips, 528
peanut insects, 69
raspberry root borer, 287
rice weevil, 64
southern potato wireworm, 879
sweet potato insects in soil, 768
tobacco wireworm, 345
tomato russet mite, 769
Typhlodromus fallacis, 513
wireworms, 66, 480
Chlorobenzilate, for control of
fowl tick, 714
two-spotted spider mite, 151, 243
winter grain mite, 519
Chlorobenzilate, toxicity of, to *Typhlodromus fallacis*, 512
Chlorthion, for control of
Acontia dacia, 765
body louse, 747
boll weevil, 713, 759, 784
bollworm, 713, 784
clover leaf weevils, 543
cotton aphid, 713
cotton insects, 73
cotton leafworm, 707
desert spider mite, 860
grain mite, 813

green June beetle, 829
green peach aphid, 25, 604
house fly, 27, 78, 622, 705, 752
northern fowl mite, 348
onion thrips, 334
peanut insects, 69
poultry lice, 349
southern garden leafhopper, 358
strawberry spider mite, 861
Cholinesterase activity, relationship of chemical structure to, 545
toxicity to, 545
Cholinesterase, inhibition of, by chlorinated phosphates, 545
malathion and isomalathion, 484
Cholinesterase in eggs of insects and its relationship to action of parathion, 777
Choristoneura
fumiferana, 338, 412, 556
pinus, 412
Chortopaga viridifasciata-viridifasciata, 371
Chrysobothris sp., 664
Chrysomyza demandata, 619
Chrysopa
oculata, 698
plorabunda, 698
rufilabris, 698
spp., 466
Chrysopidae, 199, 384
Chrysops
discalis, 87
sp., 384
Cimbicidae, 701
Circulifer tenellus, 95, 123, 388
Cixius nervosus, 203
Cloanthus frontalis, 371
Cnephasia
longana, 204
spp., 205
Cnephia pecuarum, 121
Coenosia tigrina, 336
Coenzyme A content, effect of iodoacetic acid on, 554
Colaspis flavida, 371
Coleoptera
associated with oak wilt-infected trees, 222
in or near roots of cultivated crucifers, 336
reared from oak wilt trees, 224
winter survey of in feed and flour mills, 154
Colias philodice eurytheme, 327, 703
Colladonus
clitellarius, 382
geminatus, 140, 272
Collembola, 703
Colopterus
hemipterus, 228
morio, 228
semitectus, 222
Colorado 9, for control of beet leafhopper, 124
Compounds, 3,4-methylenedioxyphenoxy, as synergists for allethrin and pyrethrins against house flies, 419
Compound 876, for control of two-spotted spider mite, 243
Conoderus
amplicollis, 880
falli, 879
sp., 69, 112, 766
vagus, 879
vespertinus, 256, 345
Conotrachelus nenuphar, 14, 198, 382, 522
Copper naphthenate, for control of house fly larvae, 75
Copper sulfate, tribasic, toxicity of, to *Typhlodromus fallacis*, 512
Copromyza annulus, 599
Corrodentia, 219
Corticaria serrata, 269
Cosmopolites sordidus, 297
Cotinis nitida, 828, 868
Cotton, effect of pink bollworm on yield and quality of, 852
Crabro, 203
Crabronine, 203
Crambus hortuellus, 205
Cremastus sp., 384
Criocerus
asparagi, 384
duodecimpunctata, 384
Crossocerus quadrimaculatus, 203
Cryolite, for control of
tobacco hornworm, 819
tomato russet mite, 769
Cryolite, effect of on
Amazon fly, 268
Cuban fly, 268
Cryptarcha ample, 226
Cryptophagus sp., 337
Cryptorhynchus pumilus, 384
Crypturgus sp., 664
Ctenicera sp., 384
Culex
annulirostris, 787
pipiens, 203, 273
quinquefasciatus, 261, 787
restuans, 273
sitiens, 787
territans, 273
tritaeniorhynchus, 261
Curculio
auriger, 717
proboscideus, 717
Curly top, 95
Curly-top virus, reservoirs of, 388
Cyclethrin, effectiveness with synergists against the body louse, 643
Cylas formicarius elegantulus, 766
Cynomyopsis cadaverina, 597
Cyrtepistomus castaneus, 717
Cytochrome oxidase, effects of insecticidal poisoning on, 675

Dacus
ciliatus, 177
cucurbitae, 134, 176
dorsalis, 134, 135, 176, 716
emmerezi, 177
melanotus, 177
passiflorae, 177
spp., 177
tryoni, 177
D-D mixture, for control of
green June beetle larvae, 868
pink bollworm in the soil, 718
wireworms, 480
DDT, breakdown of granulated formulations, 414
DDT, distribution and metabolism of in American cockroach, 43
DDT, effect of on
mites of pecan, 404
predaceous mites, 196
yields of red clover, 372
DDT for control of
Adistemia watsoni, 270
apple aphid, 522
apple leaf miner, 218
apple pests, 523
armyworms, 403
beet leafhopper, 124
boll weevil, 784, 874
bollworm, 784, 874
cabbage looper, 386
cattle lice, 237
chinch bug, 84
codling moth, 197, 468, 867
common malaria mosquito, 523
corn earworm, 540, 823
cotton insects, 72
cucumber beetles, 131
cutworms, 403
Drosophila, 609
dusky sap beetle, 540
eggplant lace bug, 423
European corn borer, 115, 411, 835
grape mealybug, 468
green June beetle, 829, 868
green peach aphid, 25
hairy spider beetle, 280
harlequin bug, 423
horn fly, 393, 396, 407
house fly larvae, 75
khapra beetle, 491
leaf miner, 468
lygus bugs, 94, 693
madrone thrips on apple, 398
northern fowl mite, 348
onion thrips, 333, 528
pea aphid, 403
peanut insects, 69
Piesma cinerea, 561
pine needle miner, 426
pine spittlebug, 426
pink bollworm, 560, 874
poultry lice, 349
potato flea beetle, 530, 558
Psiloptera drummondi, 391
raspberry root borer, 287
rosy apple aphid, 522
southern garden leafhopper, 357
southern potato wireworm, 880
spruce budworm, 338
stable fly, 396
stored corn insects, 755
Tabanus larvae, 402
three-cornered alfalfa hopper, 268
thrips on peanuts, 632
tomato russet mite, 770
woolly apple aphid, 522
DDT
joint action of with ethyl alcohol against Anopheline larvae, 433
physical changes after storage, 711
resistance of lygus bugs to, 94
DDT, radioactive, metabolism of, by
European corn borer, 579
Madeira roach, 579
DDT, toxicity of to
American cockroach, 167
bollworm, 368
Chrysopa sp., 360
corn earworm larvae, 409
Geocoris sp., 360
Hippodamia sp., 360
Nabis sp., 360
Orius sp., 360
tobacco budworm, 368, 409
Typhlodromus fallacis, 513
DDVP, for control of
boll weevil, 713, 784, 874
bollworm, 713, 784, 874
cotton aphid, 713
DDVP, formed from Bayer L 13/59, 801
DDVP, P^{32}-labeled, absorption and translocation of in a cockroach, 808

Deltocephalus flavicosta, 371
Dermestes caninus, 119
Demeton, for control of
chinch bug, 84
cotton aphid, 707
desert spider mite, 860
green peach aphid, 26, 165
insects attacking vetch, 666
peanut insects, 69
southern garden leafhopper, 357
strawberry spider mite, 860
two-spotted spider mite, 243
Demeton
methyl analog of, for control of two-spotted spider mite, 243
resistance of house fly larvae to, 463
Demeton, toxicity of to
corn planthopper, 58
Cyrtorhinus mundulus, 58
Geocoris sp., 361
Hippodamia sp., 361
Nabis sp., 361
Orius sp., 361
Dendroctonus
engelmanni, 211
pseudotsugae, 426
Deodorized APCO 467, 415
Dermanyssus gallinae, 629
Derris, variation in effectiveness against pea aphid, 363
Desmometopa sordida, 621
Diabrotica
balteata, 766
longicornis, 709
undecimpunctata howardi, 68, 131, 384, 632, 709
virgifera, 709
Diaphania
hyalinata, 872
nitidalis, 870
Diaspis carueli, 260
Diatraea saccharalis, 267
Diazinon, effect of piperonyl butoxide on anticholinesterase activities on house fly ChE, 591
Diazinon, for control of
Acontia dacia, 765
apple leaf miner, 218
apple pests, 523
body louse, 747
cabbage looper, 386
codling moth, 468
corn earworm, 540
dusky sap beetle, 540
flies in Wyoming, 343
fowl tick, 714
grape mealybug, 468
green June beetle, 829
green peach aphid, 604
horn fly, 27, 406
house fly, 27, 75, 78, 622, 704
leaf miner, 468
onion thrips, 334
potato flea beetle, 531
southern garden leafhopper, 358
stable fly, 28
two-spotted spider mite, 243
Diazinon, resistance of house fly larvae to, 463
Dichlone, for control of apple pests, 522
Dieldrin
effect of on germination and emergence of sugar beets, 710
physical changes after storage, 711
Dieldrin, for control of
Adistemia watsoni, 270
alfalfa weevil, 687
apple leaf miner, 218
apple pests, 523
billbugs on corn, 234
boll weevil, 697, 713, 784, 874
bollworm, 713, 784, 874
cabbage looper, 386
chinch bug, 84
clover leaf weevils, 543
clover root borer, 162, 641
cotton aphid, 713
cotton insects, 72
cucumber beetles, 131
European apple sawfly, 522
European corn borer, 835
green June beetle, 829, 869
horn fly, 393
house fly larvae, 75, 705
meadow spittlebug, 162
onion thrips, 334
pale western cutworm, 146
pea aphid, 637
peach tree borer, 574
peanut insects, 70
Piesma cinerea, 561
pine needle miner, 426
pine spittlebug, 426
pink bollworm, 560, 874
plum curculio, 14, 522
potato flea beetle, 531, 557
rice weevil, 64
southern garden leafhopper, 357
southern potato wireworm, 879
sweet potato insects in soil, 767
Tabanus larvae, 402
thrips on peanuts, 632
tobacco wireworm, 345
tomato russet mite, 770
wireworms, 66, 480
Dieldrin, pasture treatment, 136
Dieldrin, toxicity of to
boll weevil, 758
Chrysopa sp., 360
Geocoris sp., 360
Hippodamia sp., 360
Nabis sp., 360
Orius sp., 360
Typhlodromus fallacis, 513
Dilan, for control of
apple pests, 522
cucumber beetles, 132
Drosophila, 417
green June beetle larvae, 869
potato flea beetle, 531
Dinitroanisole, for control of body louse, 749
Diprion similis, effect of different hosts on biology of, 400
Diptera, 219, 384, 413
in or near roots of cultivated crucifers, 336
Dipterex (Bayer L 13/59) for control of green peach aphid, 25
Dirhinus
auratus, 177
giffardii, 176
Discaelius merula, 204
Disonycha pennsylvanica, 384
Ditolyl trichloroethane, for control of cucumber beetles, 132
DMC, for control of two-spotted spider mite, 243
DMC, toxicity of to
house cricket, 262
Typhlodromus fallacis, 513
Dolichopus
bifractus, 336
sp., 600
Dow ET-14
effect of on germination and emergence of sugar beets, 710
for control of house fly, 27
Dow ET-15, effect of on germination and emergence of sugar beets, 710
Draeculacephala
antica, 371
portola, 384
Drosophila
affinis, 418
algonquin, 418
buscki, 418, 608, 621
funebris, 418
hydei, 418, 608, 621
immigrans, 418, 608
melanogaster, 300, 417, 418, 429, 607, 621, 638, 708
quinaria, 418
repleta, 418
simulans, 607
sp., 703
transversa, 418
tripunctata, 418, 607
Drosophila melanogaster, length of larval period related to DDT-resistance, 671
Dry ice as an insect anesthetic, 264

Echinolaelaps echidninus, 702
Ecological studies of
beet leafhopper, 95
spider mites on cotton, 424
thrips on cotton, 423
Ectemnius rubicola, 204
Ectopsocus pumilis, 119
Elaphidion parallelum, 384
Elasmopalpus lignosellus, 68
Empoasca
fabae, 356, 371, 571
maligna, 201
solana, 356
Endoconidiophora fagacearum, 219
insect transmission of, 560
interval between inoculation and appearance of foliar symptoms, 429
Endria inimica, 371
Endrin, for control of
Acontia dacia, 765
Endrin, for the control of
boll weevil, 713, 784, 874
bollworm, 713, 784, 874
cabbage looper, 386
clover leaf weevils, 543
clover root borer, 641
cotton aphid, 713
cotton insects, 73
cotton leafworm, 707
cucumber beetles, 132
cyclamen mite, 392
European corn borer, 835
green June beetle, 829, 868
green peach aphid, 25, 604
house cricket, 262
house fly larvae, 75
onion thrips, 334
pale western cutworm, 146
Piesma cinerea, 561
pink bollworm, 874
potato flea beetle, 531, 557
rice weevil, 64
Tetranychus hicoriae, 404
tobacco flea beetle, 425
tobacco hornworm, 425, 819
tobacco wireworm, 345
tomato hornworm, 425
white-pine weevil, 878
wireworms, 66

Endrin, toxicity of to
boll weevil, 758
bollworm, 368, 409
Chrysopa sp., 360
cotton leafworm, 759
Geocoris sp., 360
Hippodamia sp., 360
Nabis sp., 360
Orius sp., 360
tobacco budworm, 368, 409
Eomenacanthus stramineus, 629
Eotetranychus
hicoriae, 717
sexmaculatus, 624
Ephestia, 630
cautella, 119
Epilachna varivestis, 384
Epitrix
cucumeris, 373, 530, 557
fuscula, 530
hirtipennis, 425, 530
subcrinata, 530
EPN, for control of
body louse, 747
cucumber beetles, 131
European corn borer, 115, 835
plum curculio, 14
tomato russet mite, 770
wireworms, 480
EPN, resistance of house fly larvae to, 463
Epuraea spp., 228
Equipment for
applying granulated insecticides, 839
cage, lightweight, for leaf feeding insects, 421
collecting Aquatic specimens, 261
counting plate for sampling mites, 705
ethyl acetate jar for trap lights, 702
grub survey, 279
mass rearing of mites, 476
mass rearing of predatory mosquitoes, 787
rapid mite counts, 478
quantitative olfactometer, 659
spot fumigation of conveying equipment, 565
spraying alfalfa, 636
Eriosoma lanigerum, 522
Erythroneura variabilis, 421
Ethyl alcohol, joint action of with DDT against Anopheline larvae, 433
Ethylene dibromide
for control of wireworms, 480
fumigation of mangoes against West Indian fruit fly, 715
Eucoila pelleranoi, 177
Eucosma certicana, 205
Euetheola rugiceps, damaging tobacco, 567
Euplilis
claudii, 203
crassinodus, 203
Eupodidae, 516
Euschistus
spp., 382
tristigmus, 384
Exoteleia pinifoliella, 426
Explosiveness of aerosol formulations, 415
Exposure, package, to vaporized lindane, 481

Fannia
canicularis, 336, 342, 461
fuscula, 336
glaucescens, 336
Fermate, toxicity of, to
Typhlodromus fallacis, 512, 513
Ferric sulfate, for control of apple pests, 523
Ferrisiana virgata, 276
Flavor, evaluation of,
following apple sprays, 523
following wireworm treatments, 479
malathion-treated asparagus, 422
Flies, emergence of, reared from grass silage, 619
Flour beetles, damage and reproduction in wheat at three moisture levels, 244
Francilia alaskensis, 596
Frankliniella
exigua, 403, 423
fusca, 423, 623
occidentalis, 403, 423
tritici, 403, 423, 632
Fucellia ariciiformis, 596
Fumigant mixtures against,
black carpet beetle, 725
confused flour beetle, 725
granary weevil, 727
green June beetle, 868
red flour beetle, 727
rice weevil, 727
Fumigation against khapra beetle with methyl bromide, 491
ethylene dibromide, ethylene dichloride, and carbon tetrachloride, 509
Fumigation
ethylene dibromide against West Indian fruit fly in mangoes, 715
for control of tobacco wireworm, 345
spot, of conveying equipment, 565
Vapam against mites and gladiolus soil diseases, 588
use of sulfur dioxide in grain treatments, 723
Fumiport, 565
Fungicides, for control of apple pests, 521
Fusarium, 67
oxysporum, 584

Galerucella nymphaeae, 384
Gallina ciega, 296
Gargaphia solani, 422
Gasoline, for control of green June beetle larvae, 868
Gelis tenellus, 413
Geocoris sp., 695
Gliricola porcelli, 88
Glischrochilus
confluentus, 226
fasciatus, 226
sanguinolentus, 226
Glyodin, for control of
apple pests, 523
codling moth, 197
Glyodin, harmless to *Typhlodromus*, 201
Glyodin, toxicity of, to *Typhlodromus fallacis*, 513
Glyphipteryx thrasonella, 205
Glypta fumiferanae, 413
Gnathocerus cornutus, 154
Gnathotrichus materiarius, 664
Goniocotes gallinae, 628
Gorytes, 203
Graminella nigrifrons, 371
Graphium rigidum, 222
Graphocephala versuta, 384
Grapholitha molesta, 382
Graphomya maculata, 599
Grypotes puncticollis, 203
Gymnodia, 215
Gypona sp., 384

Habits, feeding, of the spotted alfalfa aphid, 589
Habrocytus phycidus, 413
Haematopinus
adventicius, 88
eurysternus, 88, 239
Halictus sp., 203
HCN fumigation against
California redscale, 534
yellow scale, 534
Helina lucorum, 599
Heliothis
virescens, 368, 408
zea, 72, 278, 361, 368, 403, 408, 493, 539, 662, 706, 712, 783, 822, 872, 874, 877
Hellula undalis, 320
Helomyza serrata, 336
Hemiptera, 219, 384
Heptachlor, effect of on
corn root growth, 230
germination and emergence of sugar beets, 710
Heptachlor, for control of
Acontia dacia, 765
alfalfa weevil, 637
billbugs on corn, 234
cabbage maggot, 355
clover leaf weevils, 543
clover root borer, 162, 641
corn rootworms, 709
cotton insects, 72
cucumber beetles, 131
Drosophila, 708
European corn borer, 115, 835
green June beetle, 829
house cricket, 262
house fly larvae, 75
meadow spittlebug, 162, 636
onion thrips, 334
pale western cutworm, 146
pea aphid, 637
peanut insects, 70
potato flea beetle, 531
rice weevil, 64
southern potato wireworm, 879
sweet potato insects in soil, 767
thief ant, 329
thrips on peanuts, 632
tobacco wireworm, 345
tomato russet mite, 770
wireworms, 66, 112
Heptachlor, toxicity of to
boll weevil, 758
Typhlodromus fallacis, 513
Hercules AC-528, for control of
Acontia dacia, 765
onion thrips, 334
strawberry spider mite, 861
Hercules 528, effect of on germination and emergence of sugar beets, 710
HETP, for control of onion thrips, 528
Hippelates
bishoppi, 342, 419
dissidens, 419
particeps, 419
plebejus, 419
Hippodamia convergens, 384
Holcomb compound 326, for control of two-spotted spider mite, 243
Homoptera, 384
Honey bee, 825
colony weight changes, 735
influence of colony size and Nosema disease on winter losses, 831
mating range of, 254
multiple mating of queens, 741
ovarian development of workers, 854

pollination of greenhouse tomatoes by, 436
resistance to American foulbrood, 470
Hoplocampa testudinea, 522
Hosts, oversummering, of the wheat curl mite, 405
House fly
Egyptian, dispersal and abundance of, 214
inhibition of development by 3,4-methylenedioxyphenol compounds, 683
Hyaloides
harti, 199
vitripennis, 199
Hydrotaea bispinosa, 596
Hylastinus obscurus, 161, 371, 640
Hylemya
antiqua, 460
brassicae, 336, 354
cilicrura, 336, 461
depressa, 336
fugax, 337
literata, 337
lupini, 248
nidicola, 337
Hylemya spp. in stems and seed balls of onions at harvest, 460
Hymenoptera, 219, 384, 413
Hypera
meles, 371, 542, 662
nigrirostris, 371, 542
postica, 384, 441, 636
punctata, 371
Hypoderma lineatum, 87, 801
Hypodynerus labiatus, 204
Hysteroneura setariae, 268, 413
Hurricanes, effect of on aphid populations, 252

Iassus lanio, 203
Icerya purchasi, 421
Idiocerus alternatus, 384
Insects
affecting vetch seed production, 403
associated with oak wilt, 219
attacking red clover, 371
control by aircraft in forest, 92
control in soil on peanuts, 68
control of in sweet potato, 766
damaging chestnut foliage, 717
effect on cotton lint and seed, 172
effect on grade yield of lumber, 664
in or near roots of cultivated crucifers, 336
in stored peanuts, 119
induced lethargy and the radiation control of, 629
transmission of beet yellows virus, 789, and yellow-net virus, 792
use of adjuvants with polyhedrosis viruses, 703
vectors of sugarcane mosaic, 413
wheat and flour, effect of microwaves on, 33
Insecticides, effect of
against pink bollworm in the soil, 718
broadcast treatments for tobacco wireworms, 344
climatic conditions on cotton insect control, 760
dosage levels on residual toxicities, 757
granulated, against the salt marsh greenhead, 401
on corn root growth, 230
on germination and emergence of sugar beets, 710
poisoning on cytochrome oxidase in cockroach, 675
repeated applications on accumulation of toxic residues, 757
seed treatment to control thief ant, 329
single treatment upon biology of house flies, 310
sunlight and rainfall on toxicity, 757
systemic on green peach aphid, 164
systemic to control vectors of sugarcane mosaic, 413
to control root maggots in rutabagas, 354
Insecticides
common names of, 141
granulated against European corn borer, 113, 834
granulated, physical properties of, 840
granulated, to control overwintering boll weevils, 698
greenhouse tests against beet leafhopper, 123
new, to control green June beetle larvae, 868
organic phosphorus, for fly control, 77
phosphorus, tests against cotton insects, 874
resistance of southern potato wireworm to, 879
sampling for residues on cabbage and celery, 612
Soap as, and spreader, 266
storage tests of water-dispersible powders, 711
systemic, chemical structure and insecticidal activity of, 645
toxicity of beneficial insects, 359
Insecticides, systemic, for control of
bloodsucking flies, 87
cattle grubs, 87
lice, 88
livestock, 86, 89
Ips
calligraphus, 664
pinus, 664
Isodrin, for control of
cabbage looper, 386
clover root borer, 162, 641
cucumber beetles, 132
European corn borer, 835
green peach aphid, 165
green June beetle, 829
meadow spittlebug, 162
rice weevil, 64
tomato russet mite, 770
Isomalathion, inhibition of cholinesterase and succinoxidase by, 487
Issus coleoptratus, 203
Itoplectis conquisitor, 413

Kalotermes minor, tolerance to desiccation, 553
Karathane, toxicity of to house cricket, 262
Khapra beetle problem in
Arizona, 508
California, 490
Kurtomathrips morrilli, 562

Laemophloeus
ferrugineus, 154
pusillus, 119, 154
turcicus, 154
Larrada semirufa, 615
Lasioderma serricorne, 119
Lathetieus oryzae, 155
Lathidius nigritus, 269
Lauseto neu, for control of northern fowl mite, 348
Lead arsenate, for control of
apple pests, 523
codling moth, 468
grape mealybug, 468
leaf miner, 468
potato flea beetle, 531
Lead arsenate, toxicity of, to *Typhlodromus fallacis*, 513
Lepidoptera, 219
Leptocera
acutangula, 621
elegans, 621
exiguella, 620
ferruginata, 621
frontinalis, 419
limosa, 521
mirabilis, 621
vagans, 342
Leptocoris trivittatus, 398
Leptostylus sp., 664
Leptothrips mali, 199
Lestica subterraneas, 204
Leucocytozoon smithi, 121
Leucophaea, 337
maderae, 579, 731
Lice, tests with systemic insecticides against, 88
Limonius
californicus, 99
canus, 99
Lindane, effect of on
grain mite, 813
yields of red clover, 372
Lindane, for control of
borers attacking sawlogs, 665
brown dog tick, 282
cattle lice, 139, 237
clover root borer, 162, 641
cotton insects, 73
cucumber bettles, 131
Drosophila, 609
eggplant lace bug, 423
green June beetle, 829, 868
green peach aphid, 25, 603
harlequin bug, 423
house cricket, 262
house fly larvae, 75
juniper scale, 260
khapra beetle, 508
meadow spittlebug, 162
northern fowl mite, 348
peanut insects, 70
pine needle miner, 426
pine spittlebug, 426
poultry lice, 349
rice weevil, 64
screw-worm, 86
silverfish, 271
stored corn insects, 753
thief ant, 329
wireworms, 66, 112, 480
Lindane
impregnated sawdust for protection of stored corn 753
metabolism of, by house flies, 548
physical changes after storage, 711
toxicity of, to *Typhlodromus fallacis*, 512
translocation of, in corn plants, 563
vapor, contamination of food and air by, 212
vaporized, package exposure to, 481
Linognathus vituli, 88, 138, 239
Liothrips urichi, 562

Lipid content of cockroach, effect on toxicity of DDT, 167
Lithocolletis
crataegella, 217
sp., 468
Lixophaga diatreae, 267
Longistigma caryae, 718
Loxostege spp., 403
Lucilia
illustris, 596
spp., 217
Lyctocoris campestris, 119
Lyctus planicollis, 127, 263
Lygus
hesperus, 94, 689
lineolaris, 94, 371, 382

Machines, different types for applying granulated insecticides, 839
Macrocentrus sp., 384
Macrocheles sp., 410
Macropsis trimaculata, 384
Macrosiphum
pisi, 363, 371, 403, 636, 771, 878
sanborni, 645
solanifolii, 252, 772
Macrosteles fascifrons, 371
Malacasoma
americanum, 556, 659
disstria, 659
fragile, 651
Maneb, toxicity of, to *Typhlodromus fallacis*, 512
Manuelia, 203
Manuscript rules and suggestions, 575
Malathion, effect of on
germination and emergence of sugar beets, 710
grain mite, 813
piperonyl butoxide on anticholinesterase activities on house fly ChE, 591
Tetranychus hicoriae, 404
yield of red clover, 374
Malathion, P^{32}-labeled, fate of in
American cockroach, 187
calves, 679
laying hen, 185
white mouse, 187
Malathion, for control of
Acontia dacia, 765
alfalfa weevil, 637
apple aphid, 522
apple leaf miner, 218
apple pests, 523
armyworms, 403
body louse, 747
brown dog tick, 282
California red scale, 103
cattle lice, 237
chicken body louse, 447, 806
chicken mite, 628
clover leaf weevils, 543
clover root borer, 641
corn earworm, 540
cotton aphid, 707
cotton leafworm, 707
cucumber beetles, 132
cutworms, 403
desert spider mite, 860
Drosophila, 417, 709
dusky sap beetle, 540
eggplant lace bug, 423
European corn borer, 115, 835
Ferrisiana virgata, 277
flies in Wyoming, 343
fowl tick, 447, 714
green June beetle, 829, 868
green peach aphid, 25, 603
harlequin bug, 423
horn fly, 407
house fly, 27, 75, 78, 307, 622, 705, 752
juniper scale, 260
khapra beetle, 509
Mallophaga sp., 628
mites in earthworm beds, 410
northern fowl mite, 348
onion thrips, 334
pea aphid, 403, 637
pine spittlebug, 426
potato flea beetle, 531
potato leafhopper, 571
poultry lice, 349
rice weevil, 64
rosy apple aphid, 522
sarcoptic mange of hogs, 140
southern garden leafhopper, 358
strawberry spider mite, 861
taxus mealybug, 274
three-cornered alfalfa hopper, 268
thrips on peanuts, 633
two-spotted spider mite, 243
winter grain mite, 519
woolly apple aphid, 522
Malathion
inhibition of cholinesterase and succinoxidase by, 484
resistance of house fly larvae to, 463
Malathion, toxicity of to
corn earworm larvae, 409
Geocoris sp., 361
Hippodamia sp., 361
Nabis sp., 361
Orius sp., 361
poultry, 447
tobacco budworm, 409
Typhlodromus fallacis, 512
Malrin, for control of house fly, 622
Megalopyge opercularis, rearing of, 37
Melanoplus
femur-rubrum, 371, 662
sp., 618
Melanotus
communis, 66
spp., 479
Melanotrichus flavosparsus, 373
Melittia cucurbitae, 872
Mellinus, 203
Menacanthus stramineus, 347, 447, 806
Menopan gallinae, 347, 628
Mesostenus gracilis, 119
Metabolites, sulfoxide and sulfone plant, of thiol isomer of Systox, 147
Metacide, for control of
cucumber beetles, 132
green peach aphid, 603
Mites in earthworm beds, 410
rusty plum aphid, 414
two-spotted spider mite, 243
winter grain mite, 519
Metagonistylum mimense, 267
Metamasius sericeus, 297
Meta-Systox, for control of peanut insects, 69
Metatetranychus
citri, 138, 624
ulmi, 197, 350, 458, 468, 522
Methocel, 703
Methods
airplane spraying for spruce budworm, 338
analysis for piperonyl butoxide by chromatography, 730
applying chemicals against root maggots in rutabagas, 350
applying insecticides against onion thrips, 529
beet yellows virus transmission by green peach aphid, 789
biological testing sprays against forest insects, 555
cattle louse control with treated rubbing devices, 236
control by aircraft, 92
controlling corn earworm without insecticides, 278
control of wireworms in sweet potatoes, 479
determination of insecticide residues in soil, 638
determining metabolism of Bayer L 13/59 in a cow, 801
determining effect and extent of lygus bug injury to carrot seed, 690
for studying mating of queen honey bees, 741
fumigating to control khapra bettle, 491, 509
fumigation against West Indian Fruit fly, 715
house fly egg-measuring, 49
impregnated paper in nutritional studies with Douglas-fir beetle, 426
labelling lone star tick with radioactive P^{32}, 393
laboratory rearing of puss caterpillar, 39
litter treatment with malathion against chicken body louse, 806
marking and trapping house flies, 214
mass rearing of pink bollworm, 559
mass rearing *Typhlodromus*, 476
mass rearing of two-spotted spider mite, 476
Oriental fruit fly, medium for mass-culture, 134
oviposition medium for Mexican fruit fly, 570
planting dates to avoid European corn borer, 877
preparation of P^{32}-labeled Bayer L 13/59 and DDVP, 808
rearing *Aedes aegypti* in chemically defined medium, 313
rearing carpet beetle, 277
rearing citrus red mite, 138
rearing cockroaches, 411
rearing fruit flies, 135
rearing pink bollworm on synthetic media, 7
rearing southern lyctus beetle, 127
sampling cabbage and celery for insecticide residues, 612
sampling drop size of aerial spray deposits, 81
sampling mite populations, 705
screening wheat varieties for resistance to the greenbug, 546
self-treating devices for cattle lice, 138
spraying alfalfa for insects, 636
spraying cloth patches with fabric protectants, 570
studying southern lyctus beetle, 263
tagging large numbers of live adult insects with radioisotopes, 210
overwintering in the soil, 718
testing grain protectants against the grain mite, 813
to study nutrition of pink bollworm, 454
transformation of counts for analysis of variance, 179

using dry ice as an insect anesthetic, 265
use of systemics on livestock, 89
use of virus against tent caterpillar, 653
Methoxychlor, effect of on
germination and emergence of sugar beets, 710
grain mite, 813
Methoxychlor, for control of
apple pests, 523
biting flies, 465
cattle lice, 237
clover leaf weevils, 543
clover root borer, 641
clover root curculio, 641
codling moth, 468
cucumber beetles, 132
Drosophila, 609
eggplant lace bug, 423
flies in Wyoming, 343
grape mealybug, 468
green June beetle, 829, 868
harlequin bug, 423
horn fly, 396, 407
leaf miner, 468
meadow spittlebug, 426
pine needle miner, 426
pine spittlebug, 426
plum curculio, 14, 522
potato leafhopper, 571
stable fly, 396
three-cornered alfalfa hopper, 268
Methoxychlor, toxicity of, to *Typhlodromus fallacis*, 513
Methyl bromide, for control of khapra beetle, 491, 509
Methyl parathion, for control of
cotton leafworm, 707
desert spider mite, 860
onion thrips, 334
strwaberry spider mite, 861
tomato russet mite, 770
3,4-Methylenedioxyphenol compounds, development in the house fly inhibited by, 683
Metriona bicolor, 384
Microbembix, 203
Microgramme
arga, 269
costulata, 269
ruficollis, 269
Microwaves, effect of on wheat and flour insects, 33
Milky disease, effectiveness of in Ohio, 4
Mimesa bicolor, 203
Mites
associated with gladiolus soil diseases, 584
counting plate for sampling populations, 705
in cricket cultures, 262
infesting earthworm beds, 409
new one on corn, 265
responses of apple trees to, 350
spider, air drift of, 624
spider, on cotton, 712
wheat curl, dry bulb or tulip bulb, 1
Mnemonica auricyania, 718
Mode of action of some chlorinated phosphates, 544
Moisture content in wheat, effect of on damage and reproduction by flour beetles, 244
Monarthrum
fasciatum, 220
mali, 220
Monochamus sp., 664
Monolexis lycti, 127
Morator nudus, 52
Mosquitoes in farm ponds, 273
Mosaic, sugarcane, insect vectors, of 413
Musca domestica, 74, 283, 307, 310, 341, 419, 453, 463, 484, 496, 544, 548, 556, 591, 597, 619, 622, 671, 683, 704, 714, 751, 820
domestica vicina, 215
sorbens sorbens, 215
Musca domestica
control in dairy barns, 27
copper content of, 569
egg-measuring techniques, 49
Muscina, 215
assimilis, 337, 419, 596
stabulans, 217, 337, 342
Murgantia histrionica, 328, 422
Myospila meditabunda, 337, 599
Myzocollis trifolii, 289
Myzus
persicae, 24, 164, 252, 602, 771, 789
solani, 137, 252

Nabam, for control of apple pests, 523
Nabis sp., 695
Naphthalene, for control of house fly larvae, 75
Nectar
extrafloral of cotton as food for pink bollworm, 875
vetch as a source of, 666
Nematocampa limbata, 718
Neoclytus acuminatus, 384
Neoleria tibialis, 596
Neotran, effect of against grain mite, 813
Neuroptera, 384
Nitidulidae, 211, 219
Nicotine sulphate
for control of apple leaf miner, 218
green peach aphid, 25
onion thrips, 528
Norvellina seminuda, 384
Nosema
disease, 831
infected bees, effect on position in the winter cluster, 743
sp., 134
NPD, for control of
cucumber beetles, 132
two-spotted spider mite, 243
NDP, resistance of house fly larvae to, 463
Nutritional medium for pink bollworm, 875

Obituary
George Adam Dean, 578
Royce Burton Knapp, 574
Leonard Marion Peairs, 430
Odynerus, 204
Oecanthus nigricornis quadripunctatus, 662
Oedaleonotus enigma, 615
Oil, for control of
corn earworm, 540
dusky sap beetle, 540
California red scale, 534
khapra beetle, 491
yellow scale, 534
Oil sprays, effect of timing on Valencia oranges, 376
Olfactometer, quantitative, 659
Oligonychus
bicolor, 718
pratensis, 265
punicae, 624
Oncometopia undata, 384
Oncopeltus fasciatus, 777
Ophiostoma pluriannulata, 222
Ophyra leucostoma, 621
Opius
concolor, 177
compensans, 176
fletcheri, 177
formosanus, 176
fullawayi, 178
humilis, 177
incisi, 176
krausii, 176
longicaudatus
var. *malaiensis*, 176
var. *novocaledonicus*, 176
var. *taiensis*, 176
melleus, 178
oophilus, 176
tryoni, 177
vandenboschi, 176
watersi, 177
Orius sp., 695
Orthotomicus caelatus, 664
Oryzaephilus surinamensis, 119, 154, 753
Oscinella
carbonaria, 419
coxendix, 419
neocoxendix, 336
Ovex, for control of
desert spider mite, 860
European red mite, 350
strawberry spider mite, 861
two-spotted spider mite, 243, 350
winter grain mite, 519
Ovex, toxicity of, to *Typhlodromus fallacis*, 513
Ovotran, toxicity of to house cricket, 262
Oxylebus, 203
Oxytelus sp., 337

Palorus
ratzeburgi, 155
subdepressus, 155
Panasol AN-1, 415
Panasol AN-3, 415
Panogen PL-1, for control of thief ant, 329
Panogen PL-2, for control of thief ant, 329
Paragopsis strigatus, 336, 461
Para-oxon, for control of body louse, 747
Paraphlepsius irroratus, 371, 384
Parascaptomyza disticha, 419
Parasites
egg, 140
from two jack-pine budworm outbreaks, 412
of *Ferrisiana virgata*, 277
of sugarcane borer, 267
Parasitus sp., 410
Paratetranychus citri, 458
Paratheresia claripalpis, 267
Parathion, for control of
Acontia dacia, 765
apple aphid, 522
apple leaf miner, 218
apple pests, 523
armyworms, 403
billbugs on corn, 234
body louse, 747
cabbage looper, 386
California red scale, 106
chinch bug, 84
clover leaf weevils, 543
clover root borer, 162
cotton aphid, 707, 761

cotton insects, 72
cotton leafworm, 707, 759
cucumber beetles, 132
cutworms, 403
Drosophila, 609
European corn borer, 835
green June beetle, 829, 868
green peach aphid, 25, 165, 603
house fly larvae, 75
khapra beetle, 491
meadow spittlebug, 162
mites in earthworm beds, 410
onion thrips, 334, 528
pea aphid, 403
peanut insects, 70
plum curculio, 14
potato flea beetle, 531
raspberry root borer, 287
rice weevil, 64
rosy apple aphid, 522
southern garden leafhopper, 357
strawberry spider mite, 861
three-cornered alfalfa hopper, 268
thrips on peanuts, 633
tomato russet mite, 770
two-spotted spider mite, 243
winter grain mite, 519
woolly apple aphid, 522
Parathion, for the eradication of
California red scale, 534
yellow scale, 534
Parathion
ovicidal action of, 777
resistance of house fly larvae to, 463
Parathion, toxicity of, to
Chrysopa sp., 360
cotton aphid, 761
desert spider mite, 762, 860
Geocoris sp., 360
Hippodamia sp., 360
Nabis, sp. 360
Orius sp., 360
two-spotted spider mite, 762
Typhlodromus fallacis, 512
Paregle
cinerella, 337
mystacea, 596
radicum, 337, 596
Particle size, effect of on European corn borer control, 836
Peach gummosis and meadow spittlebug, 382
Pectinophora gossypiella, 7, 158, 173, 454, 508, 559, 560, 562, 718, 852, 874, 875, 876
Pediculus humanus humanus, 643, 747
Pegomya ruficeps, 337
Pemphedon lethifer, 204
Penetration of packaging materials by lindane, 483
Penicillium, 67
Pentachlorocyclohexene, an intermediate in the metabolism of lindane, 548
Penthaleus major, 515
Peregrinus maidis, 59
Perezia mesnili, 320
Peridroma margaritosa, 403
Periplaneta americana, 166, 484, 554, 555, 662, 675, 808, 862
Periplaneta americana,
differences in arsenic tolerance, 39
distribution and metabolism of DDT in, 43
Perthane, for control of
beet leafhopper, 124
clover root borer, 162
cucumber beetles, 132
eggplant lace bug, 423
harlequin bug, 423
horn fly, 407
meadow spittlebug, 162
onion thrips, 334
potato leafhopper, 571
southern garden leafhopper, 357
wireworms in sweet potatoes, 480
Pesticides, selective, as aids to biological control of apple pests, 196
Petrobia
apicalis, 291
harti, 745
latens, 745
Phaeogenes
hariolus, 413
sericata, 342
Phaonia errans, 599
Phenothiazine, for control of house fly larvae, 75
Philaenus leucophthalmus, 161, 371, 382, 636
Philanthus triangulum, 203
Phorocera sp., probably *claripennis*, 39
Phosphates, chlorinated, relative toxicity and mode of action of, 544
Phosphorus
compounds against body lice, 747
radioactive, fate of in house flies and cockroaches, 820
Phryxe pecosensis, 413
Phyllophaga spp., 766
Phyllotreta striolata, 337
Phymatodes amoenus, 384
Phytonomus, 204
Phytophaga destructor, feeding mechanism of, 182
Pieris rapae, 57, 320, 421
Piesma cinerea, 561
Piophila
affinis, 599
vulgaris, 599
Piperonyl butoxide, effect of, on
action of malathion on house fly strains, 307
anticholinesterase activities of phosphorus insecticides on house fly ChE, 591
grain mite, 813
purified bovine erythrocyte ChE, 591
Piperonyl butoxide, for control of
Drosophila, 417, 609
Piperonyl butoxide, C^{14}-labeled synthesis of, and its fate in the Madeira roach, 729
Pirazinon, for control of house fly, 78
Pissodes strobi, 211, 556, 664, 878
Plathypena scabra, 403
Pleocoma
crinita, 431
oregonensis, 431
Plodia interpunctella, 119
Plutella maculipennis, 320, 386
Pollenia rudis, 337, 621
Pollination of
apples with bee-collected pollen, 825
greenhouse tomatoes by honey bees, 436
hairy vetch by honey bees, 666
white Dutch clover, factors affecting, 685
Polygraphus rufipennis, 664
Popillia japonica, 4, 327, 371, 566, 717
Population surveys of corn earworm in Central Florida, 493
Potasan, for control of body louse, 747
Predators
Chrysopid, of the cotton aphid, 698
hymenopterus, as control agents, 202
Proecidochares utilis, 178
Prodenia
eridania, 239
ornithogalli, 421
Prolan, for control of cucumber beetles, 132
Protoparce
quinquemaculata, 425
sexta, 425, 818
Protophormia terrae-novae, 596
Psallus seriatus, 783
Psen ater, 203
Pseudococcus
boninsis, 268, 414
cuspidatae, 273
maritimus, 468
Pseudaletia unipuncta, 52, 384, 403
Pseudomonas marginata, 584
Pseudophycus meritorius, 277
Pseudopityophthorus
minutissimus, 560
pruinosus, 220, 561
Psiloptera drummondi, 391
Psorophora
ciliata, 416
confinis, 416
Psyllobora vigintimaculata, 384
Pterocallidium sp., 289, 589
Ptinus
fur, 155
hirtellus, 155
villiger, 280
Publication policies and manuscript rules, 575
Puratized agricultural spray, toxicity of, to *Typhlodromus fallacis*, 512
Pyrausta nubilalis, 19, 113, 270, 384, 410, 579, 834, 846, 877
Pyrazoxon, for control of
onion thrips, 334
southern garden leafhopper, 358
Pyrazoxon, toxicity tests for control of *Acontia dacia*, 765
Pyrenone-606, for control of *Drosophila*, 417
Pyrethrins, effectiveness with synergists against the body louse, 644
Pyrethrins, for control of *Drosophila*, 417
Pyrethrins-piperonyl butoxide, for control of
hairy spider beetle, 280
rice weevil, 64
Pyrethrum, effect of
acetone on toxicity of, 849
temperature on the action in a cockroach, 862
Pyrethrum
activators, effect of on toxicity of sabadilla to house fly, 283
an emulsifiable concentrate for, 719
effectiveness of in a thermal fog machine, 610
for control of *Drosophila*, 609
Pyroderces rileyi, 753
Pyrophosphate compounds, 647

Quarantines
against khapra beetle, 509
plant, new trends in, 881
Quedius sp., 337

Radiation
control of insects, 629
ultraviolet and visible attraction of pink bollworm moths to, 158

Radioactive materials
allethrin, 135
C^{14}-labeled piperonylbutoxide, 729
C^{14}-pyrethrum for tracer studies, 862
fate of radiophosphorus, $KH_2P^{32}O_4$ in flies and cockroaches, 820
fate of P^{32}-labeled malathion in calves, 679
P^{32}-labeled Bayer L 13/59, metabolism of, in a cow, 801
P^{32}-labeled Bayer L 13/59 and DDVP absorption and translocation in a cockroach, 808
P^{32}-labeled malathion, 185
P^{32}-labeled Systox isomers, 739
p,p'-C^{14}-ring-labeled DDT, 579
p,p'-C^{14}-ring-labeled DDE, 579
strontium, 89, 714
use of P^{32}, 393
Radioactive material, use of
$Ag^{110}NO_3$ for tagging adult insects, 210
NaI^{131} for tagging adult insects, 210
$Na_2Ir^{192}Cl$ for tagging adult insects, 210
$Rb^{86}Cl$ for tagging adult insects, 210
$Se^{46}Cl_3$ for tagging adult insects, 210
Radioisotopes, tagging large numbers of live insects with, 210
Radiostrontium, action of, in
German cockroach, 714
house fly, 714
Rahbdepyris zeae, 119
Relative humidity, effect of, on toxicity of derris to the pea aphid, 364
Repellents
butoxy polypropylene glycol against *Drosophila*, 418
pyrethrum sprays ineffective against *Drosophila*, 418
studied by olfactometer, 663
Residues, insecticide, in or on
animal tissues, 90
corn treated with DDT granules and emulsions, 846
determination in soil, 638
dieldrin, following pasture treatment, 136
effect of different impregnated carriers, 848
effect of size of granules applied, 848
effect when applied with different machines, 849
hens treated with malathion, 629
malathion-treated asparagus, 422
persistence of, in soil, 639
red clover hay, 163
sampling for on cabbage and celery, 612
Resistance, insecticide
brown dog tick to chlordane, 281
cabbage looper to DDT, 387
California red scale to HCN, 535
codling moth to DDT, 866
common malarial mosquito to DDT, 523, 567
Drosophila to DDT, 300, 671
house fly to DDT, 671
German cockroach to chlordane, 130
house flies in Japan to insecticides, 556
house flies to insecticides, 548
lygus bugs to DDT, 94
potato flea beetle to DDT, 533, 558
southern potato wireworm to insecticides, 879
transmission of between strains of mites, 858
Resistance, insecticide, comparisons of organic phosphorus against house fly larvae, 463
Resistance, plant,
alfalfa to spotted alfalfa aphid, 289
sorghums to two biotypes of corn leaf aphid, 498
wheat varieties to the greenbug, 546
Resistance to American foulbrood in honey bees, 470
Rhagium lineatum, 664
Rhagoletus
cingulata, 178
completa, 178
fausta, 178
pomonella, 178
Rhipicephalus sanguineus, 281
Rhizoglyphus
echinopus, 584
rhizophagus, 584
solani, 584
Rhopalosiphum maidis, 414, 498
maidis, 414, 498
pseudobrassicae, 698
Rhyzopertha dominica, 119, 155
Rohm & Haas FW-293, acaricidal properties of, 458
Rohm & Haas FW-293, for control of
apple leaf miner, 218
citrus red mite, 458
cyclamen mite, 392
European red mite, 458
fowl tick, 714
Pacific spider mite, 458
privet mite, 392
Schoenei spider mite, 458
two-spotted spider mite, 243, 244, 392, 458
Rohm & Haas FW-293, toxicity of, to *Typhlodromus fallacis*, 514
Rotenone, for control of
cucumber beetles, 131
European corn borer, 835
pea aphid, 364
potato leafhopper, 571
Rubbing devices, treated, to control horn flies on cattle, 407
Ryania, effect of, on
Amazon fly, 268
Cuban fly, 268
grain mite, 813
Ryania, for control of
apple leaf miner, 218
apple pests, 523
codling moth, 197, 468, 522
European corn borer, 835
European red mite, 522
grape mealybug, 468
green June beetle larvae, 868
leaf miner, 468
rice weevil, 64
two-spotted spider mite, 522
Ryania, toxicity of, to *Typhlodromus fallacis*, 512

Sabadilla, for control of
eggplant lace bug, 422
green June beetle larvae, 868
harlequin bug, 422
Sabadilla, toxicity of to house fly increased by synergists, 283
Sanninoidea exitiosa, 397, 574, 777
Sarcophaga
crassipalpus, 862
sinuata, 337
sp., 217, 596
Sarcoptes scabiei suis, 140
Scambus hispae, 413
Scaptomyza
adusta, 419, 607
graminum, 336
Scarabaeidae, 702
Scatopse fuscipes, 342, 621
Sceliphron, 204
Schizotetranychus, 745
Schradan, effect of, on cotton mites, 250
Schradan, for control of
beet leafhopper, 124
green peach aphid, 165
peanut insects, 69
rusty plum aphid, 414
two-spotted spider mite, 243
Schradan, toxicity of, to
corn planthopper, 58
Cyrtorhinus mundulus, 58
Typhlodromus fallacis, 512
Scoliocentra fraterna, 600
Scolothrips sexmaculatus, 860
Scopeuma
furcatum, 596
impudicum, 596
lanatum, 600
molle, 596
Seed Guard, for control of thief ant, 329
Sepsis sp., 419
Shell OS-2046, for control of
cabbage looper, 386
cotton aphid, 707
cotton leafworm, 707
green peach aphid, 26
Shell OS-2046, toxicity of, to
corn earworm larvae, 409
tobacco budworm larvae, 409
Simulium
congareenarum, 121
decorum, 122
jenningsi, 121
slossonae, 121
venustum, 122
Sipha flava, 268
Siphocoryne indo-brassicae, 266
Siphona irritans, 393, 395, 406, 407, 446
Sitona, 204
cylindricollis, attractiveness of sweetclover varieties to, 444
flavescens, 371
hispidula, 371, 641
Sitophylus
granarius 35, 154
oryza, 64, 119, 154, 239, 753
Sitotroga cerealella, 119, 316, 753
Sodium arsenite, for control of house fly larvae, 75
Sodium pentachlorophenate, for control of house fly larvae, 75
Solenopsis molesta, 329
Solenoptes capillatus, 138, 239
Solvents, nonexplosive mixtures of, in formulations for use in aerosol generators, 415
Solvesso-100, 415
Solvesso-150, 415
Sovacide 544C, 415
Spalangia
philippinensis, 177
sp., 177
Sphex, 203
Spissistilus festinus, 268
Spodoptera mauritia, 57
Spray deposits, drop size of aerial, 80
Sprays
combinations for control of apple pests in Connecticut, 521
synergized allethrin and pyrethrum, against mosquitoes, 156
Stauffer 1303, for control of codling moth, 470

Stegobium paniceum, 155
Steneotarsomenus pallidus, 392
Stenodynerus scabriusculus, 204
Stethorus punctum, 197
Stomoxys calcitrans, 87, 342, 395, 466, 619
Strobane, for control of
boll weevil, 613, 759, 784
bollworm, 713, 784
chinch bug, 84
cotton aphid, 713
cotton leafworm, 707
cotton insects, 73
European corn borer, 835
green peach aphid, 604
potato leafhopper, 571
rice weevil, 64
thrips on peanuts, 633
tobacco hornworm, 819
Stromatinia gladioli, 584
Succinoxidase, inhibition of, by malathion and isomalathion, 484
Sulfotepp, for control of body louse, 747
Sulphenone, for control of winter grain mite, 519
Sulfur dioxide, use of, in grain fumigants, 723
Sulfur, for control of
apple pests, 522
cotton insects, 72
desert spider mite, 860
insects attacking vetch, 666
southern garden leafhopper, 357
Tetranychus hicoriae, 404
tomato russet mite, 769
Sulphur, toxicity of to
Chrysopa sp., 360
Geocoris sp., 360
Hippodamia sp., 360
Nabis sp., 360
Orius sp., 360
Typhlodromus, 201
Typhlodromus fallacis, 513
Synanthedon pictipes, 397
Synergism with allethrin and pyrethrins, MGK-264, 156
piperonyl butoxide, 156
propyl isome, 156
sulfoxide, 156
Synergistic activity of piperonyl butoxide applied at different intervals to pyrethrum-treated house flies, 496
Synergists for pyrethrins and allethrin against house flies, 419
Symphoromyia sp., 600
Syntomosphyrum
esurus, 413
indicum, 176, 177
Syritta pipiens, 621
Syrphus sp., 461
Systox, for control of
apple leaf miner, 218
body louse, 747
corn leaf aphid, 414
rusty plum aphid, 414
winter grain mite, 519
Systox, systemic behavior of, in plants, 738
Systox, thiol isomer of, 147
Systox, toxicity of, to *Typhlodromus fallacis*, 512

Tabanus
nigrovittatus, 401
sp., 384
spp., 87, 466
Tachysphex, 203
spp., 615
tarsatus, 616
tenuipunctus, 615
Tarsostenus univittatus, 127
Tartar emetic, for control of onion thrips, 528
Taxigramma heteroneura, 618
TDE, for control of
apple leaf miner, 218
apple pests, 523
beet leafhopper, 124
Clear Lake gnat, 206
codling moth, 470
cucumber beetles, 132
Drosophila, 609
green peach aphid, 603
house fly larvae, 75
pale western cutworm, 146
tobacco flea beetle, 425
tobacco hornworm, 425, 819
tomato hornworm, 425
TDE, toxicity of to
corn earworm larvae, 409
tobacco budworm larvae, 409
Typhlodromus fallacis, 513
Temperatures, effect of on
action of malathion on house fly strains, 307
action of pyrethrum in a cockroach, 862
toxicity of DDT to cockroach, 167
toxicity of derris to the pea aphid, 364
Tenebrio
molitor, 119, 155
obscurus, 155
Tenebroides mauritanicus, 119, 154
TEPP, for control of
body louse, 747
European red mite, 350
green June beetle, 829
two-spotted spider mite, 243, 350
winter grain mite, 519
TEPP, toxicity of, to *Typhlodromus fallacis*, 514
Tetrachlorophenate, for control of house fly larvae, 75
Tetraisopropyl pyrophosphate, for control of green June beetle larvae, 868
Tetranychidae, 250, 783
Tetranychus, 197
atlanticus, 860
bimaculatus, 858
canadensis, 424
cinnabarinus, 859
desertorum, 72, 424, 760, 860
hicoriae, 404
marianae, damaging tomatoes, 712
multisetis, 859
pacificus, 458
schoenei, 198, 425, 458
telarius, 72, 151, 198, 242, 350, 392, 424, 458, 468, 476, 522, 624, 705, 745, 760, 859, 860
Tetrastichus
dacicida, 176
giffardianus, 176
Thaumatomyia trifasciata, 599
Themira putris, 621
Therioaphis
maculata, 289, 361
maculata, mechanics of feeding of, 589
Thiophosphate compounds, 646
Thiourea, for control of house fly larvae, 75
Thrips
madrone, 398
tabaci, 328, 333, 526, 562
Thyanta custator, 662
Thysanura, 271
Tortrix loeflingiana, 205
Toxaphene, effect of on *Tetranychus hicoriae*, 404
Toxaphene, for control of
Acontia dacia, 765
armyworms, 403
boll weevil, 713, 784
bollworm, 713, 784
cabbage looper, 386
cattle lice, 139
clover leaf weevils, 543
clover root borer, 641
cotton aphid, 713
cotton insects, 73
cotton leafworm, 707
cucumber beetles, 132
cutworms, 403
eggplant lace bug, 423
European corn borer, 835
green June beetle, 829, 869
harlequin bug, 423
house fly larvae, 75
insects attacking vetch, 666
lygus bugs, 94, 693
onion thrips, 334, 528
pale western cutworm, 146
pea aphid, 403
peanut insects, 69
southern garden leafhopper, 357
three-cornered alfalfa hopper, 268
thrips on peanuts, 632
tomato fruitworm, 770
tomato russet mite, 770
Toxaphene, toxicity of to
boll weevil, 758
Chrysopa sp., 360
corn earworm larvae, 409
cotton leafworm, 759
Geocoris sp., 360
Hippodamia sp., 360
Nabis sp., 360
Orius sp., 360
tobacco budworm larvae, 409
Typhlodromus fallacis, 512
Toxicity, effect of acetone on pyrethrum and allethrin, 849
Toxoptera graminum, 546
damage to wheat as affected by preceding crop, 600
eggs of, below 35th parallel, 567
Toxorhynchites
brevipalpus, 786
splendens, 786
Transmission of oak wilt fungus by bark beetle, 560
Traps for
banana weevils, 297
Egyptian house flies, 214
estimating adult densities of *Drosophila*, 417
insects associated with oak wilt, 223
insect migration studies, 383
pink bollworm moths, 158
studying movement of aphids, 771
Tribolium
castaneum, 119, 154, 244
confusum, 33, 119, 154, 244, 555, 724, 753, 816
Trichoplusia ni, 320, 385, 420
Trogoderma
granarium, 490, 508
sternalis, 119
Tubifera tenax, 621
Tychius stephensi, 371
Typhaea
fulmata, 119
stercorea, 155, 337

Typhlocyba
pomaria, 522
prunicola, 438
quercus, 438
rosae, 438
Typhlodromus
conspicuus, 196
cucumeris, 196
fallacis, 196, 476, 511
pomi, 196
Tyrophagus breviceps, 262

Ufens niger, 140
Ultraviolet radiation, attraction of pink bollworm moths to, 158

Vapam, effect of, on
gladiolus soil diseases, 584
Rhizoglyphus mites, 584
Vasates lycopersici, 768
Vectors, disease, *Simulium slossonae*, 122
Vectors, insect
banana weevils of head rot, 297
banana weevils of Panama disease, 297
Velsicol
AR-50G, 415
AR-60, 415
49-CS-56, for control of cucumber beetles, 132
Vespa
crabro, distribution and habits of, 275
orientalis, 216
Virus, curly-top, reservoirs of, 388
Virus, granulosis, to control lepidopterous pests of crucifers, 320
Virus infections
granulosis, 53
polyhedrosis, 53
Virus, polyhedral
attacking cabbage looper, 420
for control of tent caterpillar, 653
use of adjuvants with, 703
Virus, spotted wilt, not transmitted by *Kurtomathrips morrilli*, 562

Wheat streak-mosaic, oversummering volunteer wheat in the epidemiology of, 405
Willowsia platani, 745
Wireworms
control of, 65, 111
cultural practices affecting, 99
effect of treated transplanting water on control, 256
Wuchereria bancrofti, 786

Xanthomonas pruni, 382
Xylocelia tristis, 203

Zaraea americana, 701
Zarhopalus inquisitor, 277
Zethus dicomboda, 204
Zineb, for control of *Tetranychus hicoriae*, 404
Zineb, toxicity of, to *Typhlodromus fallacis*, 512
Zyzzys chilensis, 203